

### Strandings and Mortality

Marine mammals close to underwater detonations of high explosives can be killed or severely injured, and the auditory organs are especially susceptible to injury (Ketten *et al.*, 1993; Ketten, 1995). Airgun pulses are less energetic and have slower rise times, and there is no evidence that they can cause serious injury, death, or stranding. However, the association of mass strandings of beaked whales with naval exercises and, in a recent case, an LDEO seismic survey has raised the possibility that beaked whales may be especially susceptible to injury and/or stranding when exposed to strong pulsed sounds.

In March 2000, several beaked whales that had been exposed to repeated pulses from high intensity, mid-frequency military sonars stranded and died in the Providence Channel of the Bahamas Islands, and were subsequently found to have incurred cranial and ear damage (NOAA and USN, 2001). Based on post-mortem analyses, it was concluded that an acoustic event caused hemorrhages in and near the auditory region of some beaked whales. These hemorrhages occurred before death. They would not necessarily have caused death or permanent hearing damage, but could have compromised hearing and navigational ability (NOAA and USN, 2001). The researchers concluded that acoustic exposure caused this damage and triggered stranding, which resulted in overheating, cardiovascular collapse, and physiological shock that ultimately led to the death of the stranded beaked whales. During the event, five naval vessels used their AN/SQS-53C or -56 hull-mounted active sonars for a period of 16 hours. The sonars produced narrow (<100 Hz) bandwidth signals at center frequencies of 2.6 and 3.3 kHz (-53C), and 6.8 to 8.2 kHz (-56). The respective source levels were usually 235 and 223 dB re 1  $\mu$  Pa, but the -53C briefly operated at an unstated but substantially higher source level. The unusual bathymetry and constricted channel where the strandings occurred were conducive to channeling sound. This, and the extended operations by multiple sonars, apparently prevented escape of the animals to the open sea. In addition to the strandings, there are reports that beaked whales were no longer present in the Providence Channel region after the event, suggesting that other beaked whales either abandoned the area or (perhaps) died at sea (Balcomb and Claridge, 2001).

Other strandings of beaked whales associated with operation of military

sonars have also been reported (e.g., Simmonds and Lopez-Jurado, 1991; Frantzis, 1998). In these cases, it was not determined whether there were noise-induced injuries to the ears or other organs. Another stranding of beaked whales (15 whales) happened on 24–25 September 2002 in the Canary Islands, where naval maneuvers were taking place.

It is important to note that seismic pulses and mid-frequency sonar pulses are quite different. Sounds produced by the types of airgun arrays used to profile sub-sea geological structures are broadband with most of the energy below 1 kHz. Typical military mid-frequency sonars operate at frequencies of 2 to 10 kHz, generally with a relatively narrow bandwidth at any one time (though the center frequency may change over time). Because seismic and sonar sounds have considerably different characteristics and duty cycles, it is not appropriate to assume that there is a direct connection between the effects of military sonar and seismic surveys on marine mammals. However, evidence that sonar pulses can, in special circumstances, lead to hearing damage and, indirectly, mortality suggests that caution is warranted when dealing with exposure of marine mammals to any high-intensity pulsed sound.

In addition to the sonar-related strandings, there was a recent (September, 2002) stranding of two Cuvier's beaked whales in the Gulf of California (Mexico) when a seismic survey by the National Science Foundation (NSF)/LDEO vessel Ewing was underway in the general area (Malakoff, 2002). The airgun array in use during that project was the Ewing's 20-gun 8490-in<sup>3</sup> array. This might be a first indication that seismic surveys can have effects, at least on beaked whales, similar to the suspected effects of naval sonars. However, the evidence linking the Gulf of California strandings to the seismic surveys is inconclusive, and to this date is not based on any physical evidence (Hogarth, 2002; Yoder, 2002). The ship was also operating its multi-beam bathymetric sonar at the same time but, as discussed later in this document, this sonar had much less potential than these naval sonars to affect beaked whales. Although the link between the Gulf of California strandings and the seismic (plus multi-beam sonar) survey is inconclusive, this plus the various incidents involving beaked whale strandings associated with naval exercises suggests a need for caution in conducting seismic surveys in areas occupied by beaked whales.

### Non-auditory Physiological Effects

As mentioned previously, possible types of non-auditory physiological effects or injuries that might occur in marine mammals exposed to strong underwater sound might, in theory, include stress, neurological effects, bubble formation, resonance effects, and other types of organ or tissue damage. There is no proof that any of these effects occur in marine mammals exposed to sound from airgun arrays. However, there have been no direct studies of the potential for airgun pulses to elicit any of these effects. If any such effects do occur, they would probably be limited to unusual situations when animals might be exposed at close range for unusually long periods.

Long-term exposure to anthropogenic noise may have the potential to cause physiological stress that could affect the health of individual animals or their reproductive potential, which could theoretically cause effects at the population level (Gisner (ed.), 1999). However, there is essentially no information about the occurrence of noise-induced stress in marine mammals. Also, it is doubtful that any single marine mammal would be exposed to strong seismic sounds for sufficiently long that significant physiological stress would develop. This is particularly so in the case of broad-scale seismic surveys of the type planned by LDEO, where the tracklines are generally not as closely spaced as in many 3-dimensional industry surveys, or the brief acoustic measurement program planned for the northern GOM.

Gas-filled structures in marine animals have an inherent fundamental resonance frequency. If stimulated at this frequency, the ensuing resonance could cause damage to the animal. Diving marine mammals are not subject to the bends or air embolism because, unlike a human SCUBA diver, they only breath air at sea level pressure and have protective adaptations against getting the bends. There may be a possibility that high sound levels could cause bubble formation in the blood of diving mammals that in turn could cause an air embolism, tissue separation, and high, localized pressure in nervous tissue (Gisner (ed.), 1999; Houser *et al.*, 2001).

A recent workshop (Gentry (ed.), 2002) was held to discuss whether the stranding of beaked whales in the Bahamas in 2000 might have been related to air cavity resonance or bubble formation in tissues caused by exposure to noise from naval sonar. A panel of experts concluded that resonance in air-filled structures was not likely to have caused this stranding. Among other