

sandwicensis, *Wikstroemia* sp., or native ferns; and

(ii) Elevations between 319 and 959 m (856 and 3,146 ft).

Family Campanulaceae: *Cyanea koolauensis* (haha)

Oahu 20—*Cyanea koolauensis*—a, Oahu 20—*Cyanea koolauensis*—b, Oahu 35—*Cyanea koolauensis*—c, and Oahu 35—*Cyanea koolauensis*—d, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyanea koolauensis* on Oahu.

Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Slopes, stream banks, and ridge crests in wet *Metrosideros polymorpha-Dicranopteris linearis* forest or shrubland containing one or more of the following associated native plant species: *Acacia koa*, *Antidesma platyphyllum*, *Bidens* sp., *Bobea elatior*, *Broussaisia arguta*, *Cibotium* sp., *Diplopterygium pinnatum*, *Dubautia* sp., *Hedyotis* sp., *Machaerina* sp., *Melicope* sp., *Pittosporum* sp., *Pritchardia martii*, *Psychotria mariniana*, *Sadleria* sp., *Scaevola* sp., *Syzygium sandwicensis*, or *Wikstroemia* sp.; and

(ii) Elevations between 280 and 959 m (535 and 3,146 ft).

Family Campanulaceae: *Cyanea longiflora* (haha)

Oahu 4—*Cyanea longiflora*—a, Oahu 4—*Cyanea longiflora*—b, and Oahu 19—*Cyanea longiflora*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyanea longiflora* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes, bases of cliffs, or ridge crests in mesic *Acacia koa-Metrosideros polymorpha* lowland forest containing one or more of the following associated native plant species: *Antidesma* sp., *Cibotium* sp., *Coprosma* sp., *Dicranopteris linearis*, *Psychotria* sp., *Schiedea* sp., or *Syzygium sandwicensis*; and

(ii) Elevations between 146 and 1,191 m (479 and 3,906 ft).

Family Campanulaceae: *Cyanea pinnatifida* (haha)

Oahu 15—*Cyanea pinnatifida*—a, Oahu 15—*Cyanea pinnatifida*—b, and Oahu 15—*Cyanea pinnatifida*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyanea pinnatifida* on Oahu. Within these units, the

currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep, wet, rocky slopes in diverse mesic forest containing one or more of the following associated native plant species: *Canavalia* sp., *Diplazium sandwichianum*, *Pipturus albidus*, *Pisonia sandwicensis*, *Pisonia umbellifera*, *Psychotria* sp., *Strongylodon ruber*, or native ferns; and

(ii) Elevations between 450 and 881 m (1,476 and 2,890 ft).

Family Campanulaceae: *Cyanea st.-johnii* (haha)

Oahu 20—*Cyanea st.-johnii*—a and Oahu 35—*Cyanea st.-johnii*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyanea st.-johnii* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Wet, windswept slopes and ridges in *Metrosideros polymorpha* mixed lowland shrubland or *Metrosideros polymorpha-Dicranopteris linearis* lowland shrubland and containing one or more of the following associated native plant species: *Alyxia oliviformis*, *Antidesma* sp., *Bidens macrocarpa*, *Broussaisia arguta*, *Chamaesyce clusiifolia*, *Cibotium* sp., *Dubautia laxa*, *Freycinetia arborea*, *Hedyotis* sp., *Labordia* sp., *Machaerina angustifolia*, *Melicope* sp., *Psychotria* sp., *Sadleria pallida*, *Scaevola mollis*, or *Syzygium sandwicensis*; and

(ii) Elevations between 461 and 959 m (1,512 and 3,146 ft).

Family Campanulaceae: *Cyanea superba* (NCN)

Oahu 4—*Cyanea superba*—a, Oahu 4—*Cyanea superba*—b, Oahu 4—*Cyanea superba*—c, and Oahu 35—*Cyanea superba*—d, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyanea superba* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Sloping terrain on a well drained rocky substrate within mesic forest containing one or more of the following associated native plant species: *Diospyros* sp., *Hedyotis terminalis*, *Metrosideros polymorpha*, *Nestegis sandwicensis*, *Pisonia brunoniana*, *Psychotria* sp., or *Xylosma* sp.; and

(ii) Elevations between 232 and 872 m (761 and 2,991 ft).

Family Campanulaceae: *Cyanea truncata* (haha)

Oahu 20—*Cyanea truncata*—a and Oahu 21—*Cyanea truncata*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyanea truncata* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Windward slopes and stream banks in mesic to wet forests containing one or more of the following associated native plant species: *Cibotium chamissoi*, *Cyrtandra calpidicarpa*, *Cyrtandra laxiflora*, *Cyrtandra propinqua*, *Diospyros sandwicensis*, *Hibiscus arnottianus*, *Metrosideros polymorpha*, *Neraudia melastomifolia*, *Pipturus albidus*, or *Pisonia umbellifera*; and

(ii) Elevations between 54 and 705 m (177 and 2,312 ft).

Family Campanulaceae: *Delissea subcordata* (oha)

Oahu 4—*Delissea subcordata*—a, Oahu 15—*Delissea subcordata*—b, Oahu 15—*Delissea subcordata*—c, Oahu 15—*Delissea subcordata*—d, Oahu 35—*Delissea subcordata*—e, and Oahu 35—*Delissea subcordata*—f, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Delissea subcordata* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Moderate to steep gulch slopes in mixed mesic forests containing one or more of the following associated native plant species: *Acacia koa*, *Alyxia oliviformis*, *Antidesma* sp., *Bobea* sp., *Claoxylon sandwicense*, *Chamaesyce multiformis*, *Charpentiera obovata*, *Diospyros hillebrandii*, *Diospyros sandwicensis*, *Hedyotis acuminata*, *Metrosideros polymorpha*, *Myrsine lanaiensis*, *Nestegis sandwicensis*, *Pisonia* sp., *Pouteria sandwicensis*, *Psychotria hathewayi*, *Psydrax odorata*, or *Streblus pendulinus*; and

(ii) Elevations between 179 and 928 m (587 and 3,044 ft).

Family Campanulaceae: *Lobelia gaudichaudii* ssp. *koolauensis* (NCN)

Oahu 20—*Lobelia gaudichaudii* ssp. *koolauensis*—a, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Lobelia gaudichaudii* ssp. *koolauensis* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not