

limited to, the habitat components provided by:

(i) Moderate to steep slopes in *Metrosideros polymorpha* lowland wet shrublands or bogs and containing one or more of the following associated native plant species: *Bidens* sp., *Broussaisia arguta*, *Cibotium* sp., *Dicanthelium koolauense*, *Isachne distichophylla*, *Machaerina angustifolia*, *Melicope* sp., *Sadleria pallida*, *Scaevola* sp., or *Vaccinium dentatum*; and

(ii) Elevations between 383 and 867 m (1,256 and 2,844 ft).

**Family Campanulaceae: *Lobelia monostachya* (NCN)**

Oahu 30—*Lobelia monostachya*—a, Oahu 22—*Lobelia monostachya*—b, Oahu 33—*Lobelia monostachya*—c, and Oahu 35—*Lobelia monostachya*—d, identified in the legal descriptions in paragraph (i) of this section, constitutes critical habitat for *Lobelia monostachya* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep, sparsely vegetated cliffs in mesic shrubland containing one or more of the following associated native plant species: *Artemisia australis*, *Carex meyenii*, *Eragrostis* sp., or *Psilotum nudum*; and

(ii) Elevation between 79 and 592 m (259 to 1,942 ft).

**Family Campanulaceae: *Lobelia niihauensis* (NCN)**

Oahu 4—*Lobelia niihauensis*—a and Oahu 17—*Lobelia niihauensis*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Lobelia niihauensis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Exposed mesic or dry cliffs or ledges and containing one or more of the following associated native plant species: *Artemisia* sp., *Bidens* sp., *Carex meyenii*, *Dodonaea viscosa*, *Doryopteris* sp., *Eragrostis* sp., *Leptecophylla tameiameia*, *Melanthera tenuis*, *Osteomeles anthyllidifolia*, *Plectranthus parviflorus*, *Schiedea mannii*, or *Sida fallax*; and

(ii) Elevations between 407 to 926 m (1,335 to 3,037 ft).

**Family Campanulaceae: *Lobelia oahuensis* (NCN)**

Oahu 20—*Lobelia oahuensis*—a and Oahu 35—*Lobelia oahuensis*—b, identified in the legal descriptions in paragraph (i) of this section, constitute

critical habitat for *Lobelia oahuensis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes on summit cliffs in cloudswet wet forests or in lowland wet shrubland that are frequently exposed to heavy wind and rain and containing one or more of the following associated native plant species: *Bidens* sp., *Broussaisia arguta*, *Cheirodendron trigynum*, *Cibotium* sp., *Dicranopteris linearis*, *Dubautia laxa*, *Freycinetia arborea*, *Hedyotis* sp., *Labordia hosakana*, *Lycopodium* sp., *Machaerina angustifolia*, *Melicope* sp., *Metrosideros polymorpha*, *Peperomia* sp., *Phyllostegia* sp., *Sadleria squarrosa*, *Scaevola* sp., *Syzygium sandwicensis*, *Vaccinium* sp., or *Wikstroemia* sp.; and

(ii) Elevations between 415 and 959 m (1,361 and 3,146 ft).

**Family Campanulaceae: *Trematolobelia singularis* (NCN)**

Oahu 20—*Trematolobelia singularis*—a, Oahu 20—*Trematolobelia singularis*—b, Oahu 34—*Trematolobelia singularis*—c, Oahu 35—*Trematolobelia singularis*—d, and Oahu 35—*Trematolobelia singularis*—e, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Trematolobelia singularis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep, windswept cliff faces or slopes in *Metrosideros polymorpha*—*Dicranopteris linearis* lowland wet shrubland and containing one or more of the following associated native plant species: *Broussaisia arguta*, *Cibotium* sp., *Dubautia laxa*, *Eugenia* sp., *Melicope* sp., *Sadleria* sp., or *Wikstroemia* sp.; and

(ii) Elevations between 545 and 953 m (1,788 and 3,126 ft).

**Family Caryophyllaceae: *Alsinidendron obovatum* (NCN)**

Oahu 4—*Alsinidendron obovatum*—a, Oahu 4—*Alsinidendron obovatum*—b, and Oahu 15—*Alsinidendron obovatum*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Alsinidendron obovatum* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Ridges and slopes in lowland diverse mesic forest dominated by *Acacia koa* and *Metrosideros polymorpha* and containing one or more

of the following associated native plant species: *Alyxia oliviformis*, *Antidesma platyphyllum*, *Bidens torta*, *Cibotium chamissoi*, *Coprosma* sp., *Cyanea longiflora*, *Hedyotis terminalis*, *Ilex anomala*, *Machaerina* sp., *Peperomia* sp., *Perrottetia sandwicensis*, *Pipturus* sp., or *Psydrax odorata*; and

(ii) Elevations between 477 and 943 m (1,565 and 3,093 ft).

**Family Caryophyllaceae: *Alsinidendron trinerve* (NCN)**

Oahu 4—*Alsinidendron trinerve*—a, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Alsinidendron trinerve* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Slopes in wet forest or the wetter portions of diverse mesic forest dominated by *Metrosideros polymorpha* or *Ilex anomala* and *Metrosideros polymorpha* montane wet forest and containing one or more of the following associated native plant species:

*Broussaisia arguta*, *Coprosma ochracea*, *Diplazium sandwichianum*, *Gunnera* sp., *Hedyotis* sp., *Machaerina* sp., *Nothoperanema rubiginosa*, *Peperomia* sp., *Perrottetia sandwicensis*, *Phyllostegia* sp., *Pipturus albidus*, or *Vaccinium* sp.; and

(ii) Elevations between 833 and 1,233 m (2,732 and 4,044 ft).

**Family Caryophyllaceae: *Schiedea hookeri* (NCN)**

Oahu 3—*Schiedea hookeri*—a, Oahu 4—*Schiedea hookeri*—b, Oahu 4—*Schiedea hookeri*—c, Oahu 4—*Schiedea hookeri*—d, Oahu 15—*Schiedea hookeri*—e, Oahu 15—*Schiedea hookeri*—f, and Oahu 15—*Schiedea hookeri*—g, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Schiedea hookeri* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Slopes, cliffs or cliff bases, rock walls, or ledges in diverse mesic or dry lowland forest often dominated by *Metrosideros polymorpha*, *Diospyros sandwicensis*, or *Diospyros hillebrandii*, and containing one or more of the following associated native plant species: *Acacia koa*, *Alyxia oliviformis*, *Antidesma pulvinatum*, *Artemisia australis*, *Bidens torta*, *Carex meyenii*, *Carex wahuensis*, *Charpentiera tomentosa*, *Dodonaea viscosa*, *Elaeocarpus bifidus*, *Eragrostis grandis*, *Hibiscus* sp., *Leptecophylla*