

tameiameiae, *Melanthera tenuis*, *Pisonia sandwicensis*, *Pouteria sandwicensis*, *Psydrax odorata*, *Sida fallax*, or *Stenogyne* sp.; and

(ii) Elevations between 238 and 978 m (781 and 3,208 ft).

Family Caryophyllaceae: *Schiedea kaalae* (maolioli)

Oahu 4—*Schiedea kaalae*—a, Oahu 15—*Schiedea kaalae*—b, Oahu 15—*Schiedea kaalae*—c, Oahu 15—*Schiedea kaalae*—d, Oahu 20—*Schiedea kaalae*—e, and Oahu 21—*Schiedea kaalae*—f, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Schiedea kaalae* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes, cliffs, stream banks, or deep shade in diverse mesic or wet forests and containing one or more of the following associated native plant species: *Alyxia oliviformis*, *Boehmeria grandis*, *Charpentiera* sp., *Claoxylon sandwicense*, *Cyrtandra calpidicarpa*, *Cyrtandra laxiflora*, *Diospyros hillebrandii*, *Diplazium arnottii*, *Diplazium sandwichianum*, *Dryopteris unidentata*, *Freycinetia arborea*, *Hedyotis acuminata*, *Nothocestrum longifolium*, *Pipturus albidus*, *Pisonia sandwicensis*, *Pisonia umbellifera*, *Pouteria sandwicensis*, *Psychotria hathewayi*, *Selaginella arbuscula*, or *Xylosma hawaiiense*; and

(ii) Elevations between 64 and 904 m (210 and 2,965 ft).

Family Caryophyllaceae: *Schiedea kealiae* (maolioli)

Oahu 1—*Schiedea kealiae*—a, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Schiedea kealiae* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes or cliff faces or bases in dry remnant *Erythrina sandwicensis* forest and containing one or more of the following associated native plant species: *Bidens* sp., *Hibiscus arnottianus*, *Lepidium bidentatum*, *Melanthera remyi*, *Myoporum sandwicense*, *Plumbago zeylanica*, *Psydrax odorata*, *Sicyos* sp., or *Sida fallax*; and

(ii) Elevations between 47 and 341 m (154 and 1,118 ft)

Family Caryophyllaceae: *Schiedea nuttallii* (NCN)

Oahu 4—*Schiedea nuttallii*—a, Oahu 15—*Schiedea nuttallii*—b, and Oahu 15—*Schiedea nuttallii*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Schiedea nuttallii* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Rock walls, forested slopes, or steep walls in *Acacia koa*-*Metrosideros polymorpha* lowland mesic forest or *Metrosideros polymorpha*-*Dodonaea viscosa* forest and containing one or more of the following associated native plant species: *Alyxia oliviformis*, *Antidesma platyphyllum*, *Bidens torta*, *Cibotium chamissoi*, *Coprosma* sp., *Cyanea longiflora*, *Hedyotis terminalis*, *Ilex anomala*, *Machaerina* sp., *Peperomia* sp., *Perrottetia sandwicensis*, *Pipturus* sp., or *Psydrax odorata*; and

(ii) Elevations between 408 and 1072 m (1,338 and 3,516 ft).

Family Caryophyllaceae: *Silene lanceolata* (NCN)

Oahu 4—*Silene lanceolata*—a, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Silene lanceolata* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Cliff faces or ledges of gullies in dry to mesic shrubland or cliff communities and containing one or more of the following associated native plant species: *Artemisia australis*, *Bidens* sp., *Carex* sp., *Chamaesyce* sp., *Dodonaea viscosa*, *Lysimachia* sp., *Osteomeles anthyllidifolia*, *Schiedea mannii*, or *Tetramolopium filiforme*; and

(ii) Elevations between 328 to 978 m (1,076 to 3,208 ft).

Family Caryophyllaceae: *Silene perlmanii* (NCN)

Oahu 15—*Silene perlmanii*—a, Oahu 15—*Silene perlmanii*—b, Oahu 15—*Silene perlmanii*—c, and Oahu 15—*Silene perlmanii*—d, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Silene perlmanii* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep rocky slopes in *Acacia koa*-*Metrosideros polymorpha* lowland mesic forest; and

(ii) Elevations between 382 and 926 m (1,253 and 3,037 ft).

Family Convolvulaceae: *Bonamia menziesii* (NCN)

Oahu 2—*Bonamia menziesii*—a, Oahu 3—*Bonamia menziesii*—b, Oahu 4—*Bonamia menziesii*—c, Oahu 17—*Bonamia menziesii*—d, and Oahu 35—*Bonamia menziesii*—e, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Bonamia menziesii* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes or level ground in dry or mesic forest in open or closed canopy and containing one or more of the following associated native plant species: *Acacia koa*, *Alyxia oliviformis*, *Dianella sandwicensis*, *Diospyros sandwicensis*, *Dodonaea viscosa*, *Erythrina sandwicensis*, *Hedyotis terminalis*, *Leptecophylla tameiameiae*, *Melicope* sp., *Metrosideros polymorpha*, *Myoporum sandwicensis*, *Nestegis sandwicensis*, *Pisonia* sp., *Pittosporum* sp., *Pleomele* sp., *Pouteria sandwicensis*, *Psydrax odorata*, *Rauvolfia sandwicensis*, *Sapindus oahuensis*, *Sicyos* sp., *Sida fallax*, or *Waltheria indica*; and

(ii) Elevations between 81 and 658 m (266 and 2,158 ft).

Family Cyperaceae: *Cyperus trachysanthos* (puukaa)

Oahu 1—*Cyperus trachysanthos*—a, Oahu 28—*Cyperus trachysanthos*—b, Oahu 29—*Cyperus trachysanthos*—c, and Oahu 36—*Cyperus trachysanthos*—d, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cyperus trachysanthos* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Mud flats, wet clay soil, seasonal ponds, wet cliff seeps on seepy flats, coastal cliffs, or talus slopes containing *Hibiscus tiliaceus*; and

(ii) Elevations between 6 and 194 m (20 and 636 ft).

Family Cyperaceae: *Mariscus pennatiformis* (NCN)

Oahu 4—*Mariscus pennatiformis*—a and Oahu 4—*Mariscus pennatiformis*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Mariscus pennatiformis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat