

gaudichaudiana, *Streblus pendulinus*, *Zanthoxylum kauaense*, or native ferns; and

(ii) Elevations between 195 and 1,202 m (640 and 3,943 ft).

Family Lamiaceae: *Phyllostegia kaalaensis* (NCN)

Oahu 4—*Phyllostegia kaalaensis*—a, Oahu 4—*Phyllostegia kaalaensis*—b, Oahu 4—*Phyllostegia kaalaensis*—c, Oahu 4—*Phyllostegia kaalaensis*—d, Oahu 4—*Phyllostegia kaalaensis*—e, and Oahu 15—*Phyllostegia kaalaensis*—f, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Phyllostegia kaalaensis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Gulch slopes or bottoms or almost vertical rock faces in mesic forest or *Sapindus oahuensis* forest and containing one or more of the following associated native plant species:

Antidesma platyphyllum, *Claoxylon sandwicense*, *Diplazium sandwichianum*, *Freycinetia arborea*, *Hibiscus* sp., *Myrsine lanaiensis*, *Myrsine lessertiana*, *Neraudia melastomifolia*, *Pipturus albidus*, *Pouteria sandwicensis*, *Psychotria hathewayi*, *Streblus pendulinus*, or *Urera glabra*; and

(ii) Elevations between 248 and 878 m (813 and 2,880 ft).

Family Lamiaceae: *Phyllostegia mollis* (NCN)

Oahu 15—*Phyllostegia mollis*—a and Oahu 15—*Phyllostegia mollis*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Phyllostegia mollis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes or gulches in diverse mesic to wet forests and containing one or more of the following associated native plant species: *Acacia koa*, *Alyxia oliviformis*, *Antidesma platyphyllum*, *Carex meyenii*, *Chamaesyce multiformis*, *Claoxylon sandwicense*, *Diospyros hillebrandii*, *Dryopteris unidentata*, *Metrosideros polymorpha*, *Myrsine* sp., *Pipturus albidus*, *Pisonia umbellifera*, *Pouteria sandwicensis*, *Psychotria hathewayi*, or *Urera glabra*; and

(ii) Elevations between 519 to 928 m (1,702 to 3,044 ft).

Family Lamiaceae: *Phyllostegia parviflora* (NCN)

Oahu 20—*Phyllostegia parviflora*—d, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Phyllostegia parviflora* var. *lydgatei* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Moderate to steep slopes in mesic forest and containing one or more of the following associated native plant species: *Antidesma platyphyllum*, *Chamaesyce multiformis*, *Claoxylon sandwicense*, *Coprosma foliosa*, *Dryopteris unidentata*, *Myrsine lessertiana*, *Pipturus albidus*, *Pouteria sandwicensis*, *Selaginella arbuscula*, or *Xylosma hawaiiense*; and

(ii) Elevations between 529 to 881 m (1,800 to 2,890 ft).

Oahu 15—*Phyllostegia parviflora*—a, Oahu 15—*Phyllostegia parviflora*—b, and Oahu 15—*Phyllostegia parviflora*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Phyllostegia parviflora* var. *parviflora* on Oahu.

Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) *Metrosideros polymorpha* mixed lowland wet forest and containing one or more of the following associated native plant species: *Antidesma* sp., *Broussaisia arguta*, *Cheirodendron* sp., *Cibotium* sp., *Cyrtandra* sp., *Dicranopteris linearis*, *Melicope* sp., *Phyllostegia glabra*, *Pipturus* sp., *Pritchardia* sp., *Tetraplasandra* sp., *Touchardia latifolia*, or *Syzygium sandwicensis*; and

(ii) Elevations between 232 to 881 m (761 to 2,890 ft).

Family Lamiaceae: *Stenogyne kanehoana* (NCN)

Oahu 15—*Stenogyne kanehoana*—a and Oahu 15—*Stenogyne kanehoana*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Stenogyne kanehoana* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Lowland mesic forest and containing one or more of the following associated native plant species: *Acacia koa*, *Alyxia oliviformis*, *Bidens* sp., *Chamaesyce* sp., *Cibotium* sp., *Freycinetia arborea*, *Metrosideros polymorpha*, *Psychotria* sp., or *Scaevola* sp.; and

(ii) Elevations between 559 and 1,168 m (1,834 and 3,831 ft).

Family Loganiaceae: *Labordia cyrtandrae* (kamakahala)

Oahu 4—*Labordia cyrtandrae*—a, Oahu 20—*Labordia cyrtandrae*—b, and Oahu 20—*Labordia cyrtandrae*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Labordia cyrtandrae* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Shady gulches, slopes, or glens in mesic to wet forests and shrublands dominated by *Metrosideros polymorpha*, *Diplopterygium pinnatum*, and/or *Acacia koa* and containing one or more of the following associated native plant species: *Antidesma* sp., *Artemisia australis*, *Bidens torta*, *Boehmeria grandis*, *Broussaisia arguta*, *Chamaesyce* sp., *Coprosma* sp., *Cyrtandra* sp., *Dicranopteris linearis*, *Diplazium sandwichianum*, *Dubautia plantaginea*, *Lysimachia hillebrandii*, *Peperomia membranacea*, *Perrottetia sandwicensis*, *Phyllostegia* sp., *Pipturus albidus*, *Pouteria sandwicensis*, *Psychotria* sp., or *Rumex* sp.; and

(ii) Elevations between 232 and 1,233 m (761 and 4,044 ft).

Family Malvaceae: *Abutilon sandwicense* (NCN)

Oahu 4—*Abutilon sandwicense*—a, Oahu 4—*Abutilon sandwicense*—b, Oahu 4—*Abutilon sandwicense*—c, Oahu 15—*Abutilon sandwicense*—d, Oahu 15—*Abutilon sandwicense*—e, and Oahu 17—*Abutilon sandwicense*—f, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Abutilon sandwicense* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep slopes or gulches in dry to mesic lowland forest and containing one or more of the following associated native plant species: *Antidesma pulvinatum*, *Diospyros sandwicensis*, *Elaeocarpus bifidus*, *Eugenia reinwardtiana*, *Hibiscus arnottianus*, *Metrosideros polymorpha*, *Myrsine lanaiensis*, *Nestegis sandwicensis*, *Pipturus albidus*, *Pisonia* sp., *Pittosporum* sp., *Pleomele* sp., *Psydrax odorata*, *Rauvolfia sandwicensis*, *Reynoldsia sandwicensis*, or *Sapindus oahuensis*; and

(ii) Elevations between 215 and 725 m (705 and 2,378 ft).