

Family Malvaceae: *Hibiscus brackenridgei* (mao hau hele)

(i) Oahu 1—*Hibiscus brackenridgei*—a and Oahu 4—*Hibiscus brackenridgei*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Hibiscus brackenridgei* ssp. *mokuleianus* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(A) Slopes, cliffs, or arid ledges in lowland dry forest or shrubland and containing one or more of the following associated native plant species: *Bidens amplexans*, *Chamaesyce* sp., *Diospyros hillebrandii*, *Dodonaea viscosa*, *Doryopteris* sp., *Erythrina sandwicensis*, *Heteropogon contortus*, *Lepidium bidentatum*, *Melanthera remyi*, *Pleomele halapepe*, *Psydrax odorata*, *Reynoldsia sandwicensis*, *Sida fallax*, or *Waltheria indica*; and

(B) Elevations between 32 to 490 m (105 to 1,607 ft).

(ii) Oahu 5—*Hibiscus brackenridgei*—c, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Hibiscus brackenridgei* ssp. *molokaiana* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(A) Dry shrublands containing one or more of the following associated native plant species: *Doryopteris* sp., *Dodonaea viscosa*, *Heteropogon contortus*, *Sida fallax*, or *Waltheria indica*; and

(B) Elevations between 32 to 490 m (105 to 1,607 ft).

Family Myrsinaceae: *Myrsine juddii* (kolea)

Oahu 20—*Myrsine juddii*—a, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Myrsine juddii* on Oahu. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Ridge crests or gulch slopes in wet forests or shrublands dominated by *Metrosideros polymorpha* or a mixture of *Metrosideros polymorpha* and *Dicranopteris linearis* and containing one or more of the following associated native plant species: *Cheirodendron platyphyllum*, *Cheirodendron trigynum*, *Machaerina* sp., *Melicope clusiifolia*, *Psychotria mariniana*, or *Syzygium sandwicensis*; and

(ii) Elevations between 384 and 867 m (1,260 and 2,844 ft).

Family Myrtaceae: *Eugenia koolauensis* (nioi)

Oahu 4—*Eugenia koolauensis*—a, Oahu 19—*Eugenia koolauensis*—b, and Oahu 20—*Eugenia koolauensis*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Eugenia koolauensis* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Gentle to steep slopes or ridges in mesic or dry forests dominated by *Metrosideros polymorpha* or *Diospyros* sp. and containing one or more of the following associated native plant species: *Alyxia oliviformis*, *Bobea elatior*, *Carex meyenii*, *Dicranopteris linearis*, *Leptecophylla tameiameia*, *Myrsine lessertiana*, *Nestegis sandwicensis*, *Pleomele halapepe*, *Pouteria sandwicensis*, *Psydrax odorata*, or *Rauvolfia sandwicensis*; and

(ii) Elevations between 57 to 437 m (187 to 1,433 ft).

Family Orchidaceae: *Platanthera holochila* (NCN)

Oahu 20—*Platanthera holochila*—a and Oahu 20—*Platanthera holochila*—b, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Platanthera holochila* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) *Metrosideros polymorpha*—*Dicranopteris linearis* wet forest or *Metrosideros polymorpha* mixed shrubland and containing one or more of the following associated native plant species: *Broussaisia arguta*, *Cibotium* sp., *Clermontia* sp., *Coprosma* sp., *Dubautia* sp., *Gahnia* sp., *Leptecophylla tameiameia*, *Luzula hawaiiensis*, *Lycopodiella cernua*, *Lythrum maritimum*, *Polypodium pellucidum*, *Sadleria* sp., *Scaevola* sp., *Vaccinium reticulatum*, or *Wikstroemia* sp.; and

(ii) Elevations between 448 and 848 m (1,469 and 2,781 ft).

Family Plantaginaceae: *Plantago princeps* (laukahi kuahiwi)

(i) Oahu 20—*Plantago princeps*—d and Oahu 20—*Plantago princeps*—e, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Plantago princeps* var. *longibracteata* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(A) Sides of waterfalls or wet rock faces and containing one or more of the following associated native plant species: *Bidens* sp., *Coprosma granadensis*, *Eugenia* sp., *Lobelia gaudichaudii*, *Metrosideros rugosa*, or *Scaevola glabra*; and

(B) Elevations between 211 and 885 m (692 and 2,903 ft).

(ii) Oahu 4—*Plantago princeps*—a, Oahu 4—*Plantago princeps*—b, and Oahu 15—*Plantago princeps*—c, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Plantago princeps* var. *princeps* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(A) Slopes or ledges in *Metrosideros polymorpha* lowland mesic forests or shrublands and containing one or more of the following associated native plant species: *Artemisia australis*, *Bidens* sp., *Chamaesyce* sp., *Dubautia plantaginea*, *Eragrostis* sp., *Lysimachia* sp., *Pilea peploides*, or *Viola* sp.; and

(B) Elevations between 110 and 1,064 m (361 and 3,490 ft).

Family Poaceae: *Cenchrus agrimonoides* (kamanomano)

Oahu 4—*Cenchrus agrimonoides*—a, Oahu 4—*Cenchrus agrimonoides*—b, Oahu 15—*Cenchrus agrimonoides*—c, and Oahu 15—*Cenchrus agrimonoides*—d, identified in the legal descriptions in paragraph (i) of this section, constitute critical habitat for *Cenchrus agrimonoides* var. *agrimonioides* on Oahu. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Dry ridges, upper slopes, or ridges in lowland mixed mesic forest and containing one or more of the following associated native plant species: *Acacia koa*, *Alyxia oliviformis*, *Bobea* sp., *Carex wahuensis*, *Chamaesyce multifloris*, *Coprosma foliosa*, *Diospyros sandwicensis*, *Eragrostis variabilis*, *Gahnia beecheyi*, *Leptecophylla tameiameia*, *Metrosideros polymorpha*, *Nestegis sandwicensis*, *Psychotria* sp., or *Psydrax odorata*; and

(ii) Elevations between 357 and 874 m (1,171 and 2,867 ft).

Family Poaceae: *Eragrostis fosbergii* (Fosberg's love grass)

Oahu 4—*Eragrostis fosbergii*—a, identified in the legal description in paragraph (i) of this section, constitutes critical habitat for *Eragrostis fosbergii* on Oahu. Within this unit, the currently known primary constituent elements of