

document summarizes FDA's findings of effectiveness for certain combination drug Type B and Type C medicated feeds containing nicarbazin.

TABLE 6.—FINDINGS OF EFFECTIVENESS FOR USE OF CERTAIN DRUG COMBINATIONS CONTAINING NICARBAZIN IN POULTRY FEED

Type A article in g/ton	Type A article in g/ton	Type A article in g/ton	Indications for use	Limitations
Nicarbazin 90.8 to 181.6 (0.01 to 0.02 percent (pct))	Bacitracin methylene disalicylate 4 to 50	Roxarsone 22.7 to 45.4	Growing chickens: As an aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation.	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; do not use as a treatment for outbreaks of coccidiosis. As a sole source of organic arsenic; drug overdose or lack of water may result in leg weakness. Do not use in flushing mashers. Do not feed to laying hens in production. Discontinue medication 5 d before marketing the birds for human consumption to allow for elimination of the drug from edible tissue.
Nicarbazin 90.8 to 181.6 (0.01 to 0.02 pct)	Procaine penicillin 2.4 to 50		Growing chickens: As an aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis, and for increased rate of weight gain and improved feed efficiency.	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; do not use as a treatment for outbreaks of coccidiosis. Do not use in flushing mashers. Do not feed to chickens producing eggs for human consumption. Discontinue medication 4 d before marketing the birds for human consumption to allow for elimination of the drug from edible tissue.
Nicarbazin 90.8 to 181.6 (0.01 to 0.02 pct)	Procaine penicillin 2.4 to 50	Roxarsone 22.7 to 45.4	Growing chickens: As an aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> , and <i>E. brunetti</i>) coccidiosis, and for increased rate of weight gain, improved feed efficiency, and improved pigmentation.	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard; do not use as a treatment for outbreaks of coccidiosis. As a sole source of organic arsenic; drug overdose or lack of water may result in leg weakness. Do not use in flushing mashers. Do not feed to chickens producing eggs for human consumption. Discontinue medication 5 d before marketing the birds for human consumption to allow for elimination of the drug from edible tissue.