

Table 1b. 2003 OYs for minor rockfish by depth sub-groups (weights in metric tons).

Species	Total Catch ABC	OY (Total Catch)			Harvest Guidelines (total catch)			
		Total Catch OY	Recreational Estimate	Commercial OY for minor rockfish and HG for depth sub-groups	Limited Entry		Open Access	
					Mt	%	Mt	%
Minor Rockfish North x/	4,794	3,056	750	2,292	2,102	91.7	190	8.3
Nearshore		928	740	188				
Shelf		968	10	954				
Slope		1,160	0	1,156				
Minor Rockfish South y/	3,506	1,894	493	1,401	780	55.7	621	44.3
Nearshore		541	433	108				
Shelf		714	60	654				
Slope		639	0	639				

a/ ABC applies to the U.S. portion of the Vancouver area, except as noted under individual species.

b/ Lingcod was declared overfished on March 3, 1999. A stock assessment that included parts of Canadian waters was done in 2000 and updated for 2001. Following the assessment, lingcod was believed to be at 15 percent of its unfished biomass coastwide. The U.S. portion of the ABC for the Vancouver area was set at 44 percent of the total biomass for that area. The ABC of 841 mt was calculated using an Fmsy proxy of F45%. The total catch OY of 651 mt is based on a rebuilding plan with a 60 percent probability of rebuilding the stock to Bmsy by the year 2009 (Tmax). The total catch OY is reduced by 355 mt for the amount that is estimated to be taken by the recreational fishery, 3 mt for the amount estimated to be taken during research fishing, 4.3 mt for the amount estimated to be taken in non-groundfish fisheries, and by 5.2 mt for the amount estimated to be taken in the tribal fishery, resulting in a commercial OY of 284 mt. The open access total catch allocation is 54 mt (19 percent of the commercial OY) and the open access landed catch value is 43 mt. The limited entry total catch allocation is 230 mt and the landed catch value is 184 mt. The landed catch value is based on a discard mortality rate of 20 percent. Tribal vessels are estimated to land about 5.2 mt of lingcod in 2003, but do not have a specific allocation at this time.

c/ "Other species", these are neither common nor important to the commercial and recreational fisheries in the areas footnoted. Accordingly, Pacific cod is included in the non-commercial OY of "other fish" and rockfish species are included in either "other rockfish" or "remaining rockfish" for the areas footnoted.

d/ Pacific whiting - The most recent stock assessment was prepared in 2002, at which time the whiting stock was believed to be below 25 percent of its unfished biomass. Whiting was declared overfished on April 15, 2002 (67 FR 18117). The U.S.-Canada ABC of 235,000 mt is based on the 2002 assessment results with the application of an Fmsy