

greenwelliae, *Psychotria mariniana*, or *Sapindus oahuensis*; and

(ii) Elevations between 438 and 948 m (1,438 and 3,108 ft).

Family Asteraceae: *Lipochaeta micrantha* (nehe)

Kauai 7—*Lipochaeta micrantha*—a, and Kauai 11—*Lipochaeta micrantha*—b, identified in the legal descriptions in paragraph (a)(1) of this section, constitute critical habitat for *Lipochaeta micrantha* on Kauai. Within these units the currently known primary constituent elements of critical habitat for *Lipochaeta micrantha* are the habitat components provided by:

(i) Cliffs, ridges, stream banks, or slopes in mesic to wet mixed communities and containing one or more of the following associated native plant species: *Acacia koa*, *Antidesma* spp., *Artemisia australis*, *Bidens sandwicensis*, *Bobea* spp., *Chamaesyce celastroides* var. *hanapepensis*, *Diospyros* spp., *Dodonaea viscosa*, *Eragrostis grandis*, *Eragrostis variabilis*, *Hibiscus kokio*, *Lepidium bidentatum*, *Lobelia niihauensis*, *Melicope* spp., *Metrosideros polymorpha*, *Neraudia kauaiensis*, *Nototrichium* spp., *Pipturus* spp., *Plectranthus parviflorus*, *Pleomele aurea*, *Psydrax odorata*, *Rumex albescens*, *Sida fallax*, or *Xylosma hawaiiense*; and

(ii) Elevations between 127 and 1,090 m (418 and 3,574 ft).

Family Asteraceae: *Lipochaeta waimeaeensis* (nehe)

Kauai 13—*Lipochaeta waimeaeensis*—a, identified in the legal description in paragraph (a)(1) of this section, constitutes critical habitat for *Lipochaeta waimeaeensis* on Kauai. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Precipitous, shrub-covered gulches in diverse lowland forest and containing one or more of the following associated native plant species: *Artemisia australis*, *Chamaesyce celastroides*, *Dodonaea viscosa*, *Lipochaeta connata*, *Santalum freycinetianum*, *Schiedea spargulina*, or *Panicum* spp.; and

(ii) Elevations between 44 and 409 m (145 and 1,340 ft).

Family Asteraceae: *Remya kauaiensis* (NCN)

Kauai 11—*Remya kauaiensis*—a, Kauai 11—*Remya kauaiensis*—b, Kauai 11—*Remya kauaiensis*—c, Kauai 11—*Remya kauaiensis*—d, Kauai 11—*Remya kauaiensis*—e, and Kauai 12—*Remya kauaiensis*—f, identified in the legal descriptions in paragraph (a)(1) of this

section, constitute critical habitat for *Remya kauaiensis* on Kauai. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep, north or northeast-facing slopes in *Acacia koa*-*Metrosideros polymorpha* lowland mesic forest and containing one or more of the following associated native plant species: *Chamaesyce* spp., *Claoxylon sandwicense*, *Dianella sandwicensis*, *Diospyros* spp., *Dodonaea viscosa*, *Hedyotis terminalis*, *Melicope* spp., *Nestegis sandwicensis*, *Pouteria sandwicensis*, *Psychotria* spp., *Schiedea* spp., or *Tetraplasandra* spp.; and

(ii) Elevations between 560 and 1,249 m (1,836 and 4,097 ft).

Family Asteraceae: *Remya montgomeryi* (NCN)

Kauai 11—*Remya montgomeryi*—a, Kauai 11—*Remya montgomeryi*—b, and Kauai 11—*Remya montgomeryi*—c, identified in the legal descriptions in paragraph (a)(1) of this section, constitute critical habitat for *Remya montgomeryi* on Kauai. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Steep, north or northeast-facing slopes or cliffs in transitional wet or *Metrosideros polymorpha*-dominated mixed mesic forest and containing one or more of the following associated native plant species: *Artemisia australis*, *Bobea* spp., *Boehmeria grandis*, *Cheirodendron* spp., *Claoxylon sandwicense*, *Cyrtandra* spp., *Dubautia* spp., *Ilex anomala*, *Lepidium serra*, *Lysimachia* spp., *Myrsine linearifolia*, *Nototrichium* spp., *Pleomele aurea*, *Poa mannii*, *Sadleria* spp., *Scaevola* spp., *Stenogyne campanulata*, *Tetraplasandra* spp., or *Zanthoxylum dipetalum*; and

(ii) Elevations between 336 and 1,345 m (1,102 and 4,411 ft).

Family Asteraceae: *Wilkesia hobbdi* (dwarf iliau)

Kauai 11—*Wilkesia hobbdi*—a, identified in the legal description in paragraph (a)(1) of this section, constitutes critical habitat for *Wilkesia hobbdi* on Kauai. Within this unit, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Coastal dry cliffs or very dry ridges containing one or more of the following associated native plant species: *Artemisia australis*, *Dodonaea viscosa*, *Eragrostis variabilis*, *Hibiscus kokio* ssp.

saint johnianus, *Lipochaeta connata*, *Lobelia niihauensis*, *Myoporum sandwicense*, *Peperomia blanda*, *Peperomia* spp., *Peperomia tetraphylla*, *Peucedanum sandwicense*, *Psydrax odorata*, *Sida fallax*, *Waltheria indica*, or *Wilkesia gymnoxiphium*; and

(ii) Elevations between 12 and 685 m (40 and 2,246 ft).

Family Campanulaceae: *Brighamia insignis* (olulu)

Kauai 6—*Brighamia insignis*—a, Kauai 7—*Brighamia insignis*—b, and Kauai 11—*Brighamia insignis*—c identified in the legal descriptions in paragraph (a)(1) of this section, and Niihau 1—*Brighamia insignis*—a, identified in the legal description in paragraph (a)(2) of this section, constitute critical habitat for *Brighamia insignis* on Kauai and Niihau. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Rocky ledges with little soil or steep sea cliffs in lowland dry grasslands or shrublands with annual rainfall that is usually less than 170 cm (65 in) and containing one or more of the following native plant species: *Artemisia australis*, *Chamaesyce celastroides*, *Eragrostis variabilis*, *Heteropogon contortus*, *Hibiscus kokio*, *Hibiscus kokio* ssp. *saintjohnianus*, *Lepidium serra*, *Lipochaeta succulenta*, *Munroidendron racemosum*, or *Sida fallax*; and

(ii) Elevations between 0 and 748 m (0 and 2,453 ft).

Family Campanulaceae: *Cyanea asarifolia* (haha)

Kauai 4—*Cyanea asarifolia*—a, and Kauai 10—*Cyanea asarifolia*—b, identified in the legal descriptions in paragraph (a)(1) of this section, constitute critical habitat for *Cyanea asarifolia* on Kauai. Within these units, the currently known primary constituent elements of critical habitat include, but are not limited to, the habitat components provided by:

(i) Pockets of soil on sheer wet rock cliffs and waterfalls in lowland wet forests and containing one or more of the following native plant species: *Bidens* spp., *Dubautia plantaginea*, *Hedyotis centranthoides*, *Hedyotis elatior*, *Lysimachia filifolia*, *Machaerina angustifolia*, *Metrosideros polymorpha*, or *Panicum lineale*; and

(ii) Elevations between 182 and 1,212 m (597 and 3,976 ft).